



All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Books


Search for

Display Show Sort by Send to

All: 1 Review: 0

[About Entrez](#)[Text Version](#)[Entrez PubMed](#)[Overview](#)[Help | FAQ](#)[Tutorials](#)[New/Noteworthy](#)[E-Utilities](#)[PubMed Services](#)[Journals Database](#)[MeSH Database](#)[Single Citation Matcher](#)[Batch Citation Matcher](#)[Clinical Queries](#)[Special Queries](#)[LinkOut](#)[My NCBI](#)[Related Resources](#)[Order Documents](#)[NLM Mobile](#)[NLM Catalog](#)[NLM Gateway](#)[TOXNET](#)[Consumer Health](#)[Clinical Alerts](#)[ClinicalTrials.gov](#)[PubMed Central](#)

1: J Mol Cell Cardiol. 1999 Jan;31(1):167-78.

[Related Articles, Links](#)

F1-ATP synthase beta-subunit and cytochrome c transcriptional regulation in right ventricular hemodynamic overload and hypertrophically stimulated cardiocytes.

O'Brien TX, Schuyler GT, Rackley MS, Thompson JT.

Ralph H. Johnson Department of Veterans Affairs Medical Center, Department of Medicine, Medical University of South Carolina, Charleston 29425-2221, USA.

Cardiac hypertrophic growth secondary to hemodynamic pressure overload causes changes in energy requirements that may involve the transcriptional upregulation of oxidative phosphorylation genes. Therefore, two representative nuclear-encoded genes, the mitochondrial F1-ATP synthase beta-subunit (beta-subunit) and cytochrome c (cyt c), were examined in a feline chronic pulmonary artery banded right ventricular pressure-overload model. In the hypertrophying right ventricle, beta-subunit and cyt c mRNA levels increased after two and seven days, during the peak growth response. To examine cardiac transcriptional regulation, neonatal rat cardiac myocytes (cardiocytes) were transiently transfected with beta-subunit promoter constructs ranging from -1519 nucleotides (nt) upstream of transcription initiation as well as cyt c promoter constructs ranging from -726 nt. A full-length p1519beta-subunit/Luc construct was alpha-adrenergically inducible by 275% (+/-30%) with this activation being mapped to an enhancer region between -1519 to -1480 nt. Smaller constructs containing more proximal promoter elements were not inducible. Additionally, the full-length and enhancer deleted beta-subunit constructs were also inducible in electrically stimulated cardiocytes, suggesting a different mechanism of activation. Cyt c constructs containing known constitutive elements from -191 to -167 nt and -139 to -84 nt were responsible for the majority of the reporter activity of the full-length promoter but were not inducible in the presence of phenylephrine. Hence, we show that promoter regions containing elements common in other metabolism-related gene families are active in neonatal rat cardiocytes. Once more, we have identified a beta-subunit genomic region responsive to alpha-adrenergic and electrical stimulation.

PMID: 10072725 [PubMed - indexed for MEDLINE]

Display Show Sort by Send to

[Write to the Help Desk](#)[NCBI | NLM | NIH](#)[Department of Health & Human Services](#)[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)